

## FUSARIUM EAR ROT CAUSING FUMONISIN CONTAMINATION IN CORN

Given the wet weather as of late, we may start seeing ear rots and molds. As we get into grain harvest, I am seeing molds and rots similar to a few years ago when we experienced mycotoxin issues due to the wet fall weather. If you are seeing these issues, you're not the only one. Check out this article detailing Fusarium ear rot and the issues it may cause.

### Introduction

Fusarium ear rot is a corn disease caused by the fungi *Fusarium verticillioides*, *F. proliferatum* and *F. subglutinans*. In all three species the disease symptoms are similar, but only *F. verticillioides* and *F. proliferatum* produce a group of mycotoxins called fumonisin which at high enough levels can be toxic to animals, livestock, and humans.<sup>1</sup> Fusarium ear rot can begin during or after flowering (R1 growth stage) and is most problematic when hot and dry weather is followed by periods of high humidity. Rain before harvest may intensify the contamination of fumonisins in corn.

Due to health risks associated with mycotoxins, grain must be tested to determine fumonisin levels. Management practices can be implemented to help reduce the risk of Fusarium in future corn crops.

### Fusarium Identification

Infected kernels have a white-to-pink cottony mold which are scattered around the ear (Figure 1).

If *Fusarium* ear rot is suspected, scout fields at physiological maturity (R6 growth stage) to determine the severity.<sup>1</sup> If the ear rot is widespread, harvest as early as possible to possibly reduce the level of fumonisin contamination and ear rot damage to kernels. At harvest grain must be tested to determine the level of fumonisin as toxin levels are not synonymous with visible ear rot symptomology. Fumonisin can be detected in a kernel that does not have any damage to the seed coat (pericarp). In order to know if a cornfield is impacted by mycotoxins, testing can be completed by an elevator with testing capabilities, grain exchanges, or an approved independent laboratory. Harvested grain should be dried, cooled, and cleaned immediately after harvest, and stored apart from grain harvested from healthy fields.



**Figure 1. Fusarium infected kernels scattered around the ear.**



**Figure 2. Starburst pattern of Fusarium infected kernels.**

### Sampling and Testing Procedures

Fumonisin does not occur uniformly in bulk corn, and grain handlers should sample in several areas of a load or bin. Contact a toxicology laboratory for specific sampling, handling instructions and analysis. There are a variety of commercial laboratories and quick-test kits for mycotoxin analysis. Romer Labs ([www.romerlabs.com](http://www.romerlabs.com)) and Neogen ([www.neogen.com](http://www.neogen.com)) sell test strips for fumonisin analysis.

A farmer with contaminated grain has some options; however, contaminated grain cannot cross state lines and discounted prices are likely. Due to health risks associated with fumonisin, the United States Food and Drug Administration (FDA) has guidelines for the

# FUSARIUM EAR ROT CAUSING FUMONISIN CONTAMINATION IN CORN

maximum amounts of fumonisin allowed in livestock feed (Table 1). Horses and pigs are more susceptible to fumonisin poisoning than cattle and poultry.

If fumonisin contaminated grain is being used to make ethanol, the ethanol fermentation process increases the mycotoxin concentration levels in the distillers dried grain (DDG). Experiments have evaluated the mycotoxin concentrations in the DDGs and have reported to be approximately three times as high as the level in the grain. Molds can grow and mycotoxins can be produced in DDGs during transport, storage, and feeding at the farm which can further increase the fumonisin levels and danger when fed.<sup>2</sup>

## Corn Product Selection

Corn product selection is an important step in managing *Fusarium* outbreaks in corn. Products with insect resistance traits (above ground *Bt* protection), drought tolerance and other traits that will reduce stress during critical periods for infection. Corn products with insect resistant traits that protect the ear from corn earworm, European corn borer, and Western bean cutworm may have lower levels of ear rot as insect feeding can create additional pathways for the disease to infect the kernels. Additionally, physical traits, such as downward position ears at maturity can help reduce the potential production of mycotoxins by allowing the ear to shed water quickly.

## Management

- Crop rotation helps to minimize the carry-over of the pathogens from one year to the next.
- As *Fusarium* survives on crop residue tillage that buries the residue can help reduce the amount of inoculum for the following year crop. Tillage operations that only partially bury residue, such as chopping, disking, or chiseling, are also beneficial for pathogen reduction by helping breakdown crop residue.
- Optimize soil fertility to aid in plant health and reduce stress on the plant.
- Contact your local agronomist to select the best locally adapted corn products with proven disease tolerance or resistance.

## Steps for Dealing with Mycotoxins

If mycotoxin is detected above the guidance levels established by the Food and Drug Administration (FDA), farmers must adhere to steps established by the United States Department of Agriculture's Risk Management Agency. (<https://www.rma.usda.gov/en/Policy-and-Procedure/Loss-Adjustment-Standards---25000>) Farmers suspecting this issue should:

- Make insurance provider aware of suspected issue prior to harvest, storage or destruction of the cornfield.

**Table 1. United States FDA guidelines for fumonisins in animal feed.**

Animal	Levels in Corn & Corn By-products (ppm)	Feed Factor <sup>1</sup>	Levels in Finished Feeds (ppm)
Horse	5	0.2	1
Swine	20	0.5	10
Breeding ruminants, breeding poultry	30	0.5	15
Ruminants raised for slaughter <sup>2</sup>	60	0.5	30
Poultry	100	0.5	50
Pets	10	0.5	5

<sup>1</sup> Fraction of corn or corn by-product mixed into the total ration.

<sup>2</sup> Animals older than 3 months raised for slaughter.

Source: Summary of recommended levels for total fumonisin in feed FDA.

# FUSARIUM EAR ROT CAUSING FUMONISIN CONTAMINATION IN CORN

- Adjuster must collect samples of the Representative Sample Area (RSA) prior to the grain entering storage or the destruction of the field. Only the adjuster is approved to obtain samples from the standing crop.
- An Approved Insurance Provider (AIP)-Approved Testing Facility (i.e. laboratory) must complete analysis of these samples.<sup>5,6</sup>
- Farmers with questions regarding handling protocol should contact their insurance provider.

## Sources

<sup>1</sup> Kamle, M., Mahato, D.K., Devi, S., Lee, K.E., Kang, S. G. and Kumar, P. 2019. Fumonisin: Impact on Agriculture, Food, and Human Health and their Management Strategies. National Center for Biotechnology Information. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6628439/>

<sup>2</sup> Diaz-Royon, F. 2016. Mycotoxins are concentrated in distillers grains. Hoard's Dairyman. December 2016

<sup>3</sup> 2001 Summary of recommended levels for total fumonisin in feed <https://wayback.archive-it.org/7993/20170111170326/http://www.fda.gov/Food/FoodbornenessContaminants/NaturalToxins/ucm212900.htm>

<sup>4</sup> Long, B. 2017. Fusarium ear rot. Texas Plant Disease Diagnostic Lab [https://plantclinic.tamu.edu/calendar2018/fusarium\\_corn/](https://plantclinic.tamu.edu/calendar2018/fusarium_corn/)

<sup>5</sup> Pruitt, S. 2017. Mycotoxin levels on high plains pose threat to region's corn farmers. Texas Corn Producers. <https://www.myplainview.com/news/agriculture/article/Mycotoxin-levels-post-threat-to-High-Plains-corn-12209857.php>

<sup>6</sup> Loss adjustment manual standard handbook. United States Department of Agriculture. FCIC-25010 (11-2019) <https://www.rma.usda.gov/en/Policy-and-Procedure/Loss-Adjustment-Standards---25000>.

Web sources verified 04/23/2021.

## Legal Statement

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields.

Bayer and Bayer Cross are registered trademarks of Bayer Group. All other trademarks are the property of their respective owners. ©2021 Bayer Group. All rights reserved. 6005\_S6